

The *Drosophila* Estrogen-Related Receptor Directs a Metabolic Switch that Supports Developmental Growth

Jason M. Tennessen,¹ Keith D. Baker,^{1,2} Geanette Lam,¹ Janelle Evans,¹ and Carl S. Thummel^{1,*}

¹Department of Human Genetics, University of Utah School of Medicine, 15 North 2030 East, Room 2100, Salt Lake City, UT 84112-5330, USA

²Present address: Department of Biochemistry and Molecular Biology and the Massey Cancer Center, Virginia Commonwealth University School of Medicine, Richmond, VA 23298, USA

*Correspondence: carl.thummel@genetics.utah.edu

DOI 10.1016/j.cmet.2011.01.005

SUMMARY

Metabolism must be coordinated with development to provide the appropriate energetic needs for each stage in the life cycle. Little is known, however, about how this temporal control is achieved. Here, we show that the *Drosophila* ortholog of the estrogen-related receptor (ERR) family of nuclear receptors directs a critical metabolic transition during development. *dERR* mutants die as larvae with low ATP levels and elevated levels of circulating sugars. The expression of active *dERR* protein in mid-embryogenesis triggers a coordinate switch in gene expression that drives a metabolic program normally associated with proliferating cells, supporting the dramatic growth that occurs during larval development. This study shows that *dERR* plays a central role in carbohydrate metabolism, demonstrates that a proliferative metabolic program is used in normal developmental growth, and provides a molecular context to understand the close association between mammalian ERR family members and cancer.

INTRODUCTION

Metabolism must be tightly coupled with developmental progression, with distinct metabolic programs supporting the nutritional and energetic requirements of each stage in the life cycle. Relatively few studies, however, have addressed the molecular mechanisms that provide this temporal coordination. One of the best-characterized metabolic transitions occurs in the developing mammalian heart and is represented by a switch in substrate utilization from glucose and lactate metabolism during fetal stages to postnatal dependence on fatty acid oxidation (Lehman and Kelly, 2002). This switch is accompanied by the coordinate induction of genes involved in mitochondrial β oxidation and oxidative metabolism and mediated by the nuclear receptors PPAR α and ERR γ (Lehman and Kelly, 2002; Alaynick et al., 2007). Interestingly, this process reverts in pathological forms of cardiac hypertrophy, demonstrating that metabolic transitions can be associated with disease. In addition, the nutritional status of early developmental stages can have a profound effect on later metabolic health (Symonds et al., 2009). One of the

most dramatic manifestations of this interplay between nutrition and development is the impact of childhood obesity on the incidence of type 2 diabetes and obesity in adults. Obesity also impacts the timing of sexual maturation, linking a metabolic state to a key developmental transition (Ong, 2010). In spite of this important interplay between nutrition, metabolism, and development, little is understood about how these pathways are integrated.

Nuclear receptors are a specialized family of ligand-regulated transcription factors that play central roles in controlling development, growth, and metabolism (Chawla et al., 2001). They are defined by a conserved zinc-finger DNA-binding domain and a C-terminal ligand-binding domain (LBD) that can impart multiple regulatory functions. One subfamily of these receptors are the estrogen-related receptors (ERRs), represented by three paralogs in mammals, ERR α , ERR β , and ERR γ . Although some synthetic estrogen analogs can suppress the constitutive transcriptional activity of these receptors, they have no known naturally occurring ligands (Busch et al., 2004; Willy et al., 2004; Chao et al., 2006). Genetic studies in mice have demonstrated roles for ERR family members in mitochondrial biogenesis, oxidative phosphorylation, and lipid metabolism (for a review, see Tremblay and Giguère, 2007). Consistent with these functions, ERR α mutant mice are lean and resistant to diet-induced obesity (Luo et al., 2003). In addition, ERR α and ERR γ are essential for proper cardiac metabolism. ERR α is required for energy production in response to cardiac stress (Huss et al., 2007), whereas ERR γ directs a metabolic switch that allows the postnatal heart to metabolize fatty acids (Alaynick et al., 2007). Recent studies, however, suggest that ERRs play a broader role in metabolic homeostasis (Ao et al., 2008; Charest-Marcotte et al., 2010; Eichner et al., 2010). Moreover, all three mammalian ERRs are associated with cancer progression. ERR α is necessary for the normal growth of estrogen receptor-negative breast cancer tumor grafts (Stein et al., 2008), and elevated ERR α expression is associated with aggressive forms of breast cancer (Ariazi et al., 2002). In contrast, ERR β inhibits the progression of prostate cancer, and increased expression of ERR γ is correlated with a favorable clinical prognosis for breast cancer (Ariazi et al., 2002; Yu et al., 2008). These observations suggest that ERR family members coordinate cell growth and proliferation with metabolism. The molecular basis for this relationship, however, remains unclear.

Here, we present a functional study of the *Drosophila* ERR ortholog, *dERR*. The presence of only a single *ERR* gene in flies

eliminates the potential genetic redundancy between multiple members of the ERR subfamily, allowing us to determine its key ancestral functions. We show that *dERR* is an essential regulator of carbohydrate metabolism during larval stages. *dERR* mutants die during the second larval instar with abnormally high levels of circulating sugar and diminished concentrations of ATP and triacylglycerides (TAG). These metabolic defects result from decreased expression of genes involved in glycolysis, the pentose phosphate pathway, and other aspects of carbohydrate metabolism. These genes are coordinately induced midway through embryogenesis, in apparent direct response to the expression of activated *dERR* protein. Interestingly, this *dERR*-regulated metabolic state is ideally suited to promote larval growth by converting dietary carbohydrates into biomass and is strikingly reminiscent of the Warburg effect. Our studies reveal that an ERR family member coordinates metabolism with growth, indicate that a proliferative metabolic program is used in the context of normal development, and suggest that mammalian ERRs are associated with cancer through their ability to promote the Warburg effect.

RESULTS

dERR Is an Essential Metabolic Regulator

In an effort to examine *dERR* function, we generated two loss-of-function alleles at this locus (Figure 1A). Animals that carry a trans-heterozygous combination of these mutations, *dERR*¹/*dERR*², progress normally through embryogenesis and first-instar development, but die abruptly during the latter half of the second larval instar, with no apparent morphological defects (Figure S1A available online). *dERR* mutants, however, are severely metabolically compromised, with a 2-fold decrease in ATP levels relative to control second-instar larvae (Figure 1B). Despite these decreased levels of ATP, *dERR* mutants have elevated concentrations of the circulating sugar trehalose and normal glycogen concentrations, suggesting that they are unable to access their carbohydrate stores for energy production (Figures 1C and 1D). The high trehalose levels can be rescued by expression of wild-type *dERR* under the control of a ubiquitous GAL4 driver, indicating that the phenotype is due to a specific loss of *dERR* function (Figure S1B). TAG levels are also decreased in *dERR* mutants, suggesting that these animals have either increased fat catabolism or decreased TAG synthesis (Figure 1E). Similar metabolic defects are seen when comparing *dERR*¹/*dERR*² and *dERR*¹/*Df*(3L)*Exel6112* mutants with *dERR*¹/+ heterozygous controls, demonstrating that they do not arise from changes in genetic background and that *dERR*¹ represents a null allele for the locus (Figures S1C–S1E).

Genes Involved in Carbohydrate Metabolism Are Downregulated in *dERR* Mutants

Microarray analysis was used to examine the molecular basis for the metabolic defects in *dERR* mutants. A total of 906 genes are misregulated ≥ 1.5 -fold in *dERR* mutant second-instar larvae relative to controls. Gostat analysis of this data set revealed that there are few statistically significant gene ontology (GO) classes among the 572 upregulated genes. In contrast, the top 26 GO categories for the 334 downregulated genes all represent different aspects of carbohydrate metabolism (Figure 2A and




Figure 1. *dERR* Mutants Exhibit Metabolic Defects

(A) A schematic representation of the *dERR* locus is depicted along with the flanking genes *atg18* and *CG7979*. The *dERR*¹ lesions are shown, including an 84 bp deletion that removes the exon 2 splice acceptor (Δ) and point mutations in exons 2 and 3 (*). The *dERR*² deletion removes the entire *dERR* coding region and portions of the neighboring genes. The arrows at the 3' end of *atg18* indicate that it extends beyond what is depicted.

(B–E) *w*¹¹¹⁸ control and *dERR*¹/*dERR*² mutants (*dERR*^{-/-}) were collected as mid-second-instar larvae, and whole-animal homogenates were analyzed for concentrations of ATP (B), trehalose (C), glycogen (D), or TAG (E). Amounts of ATP, glycogen, and TAG were normalized to soluble protein levels. Mutant animals contain lower levels of ATP ($p < 1 \times 10^{-4}$) (A) and TAG ($p < 0.001$) (E) but have higher concentrations of trehalose ($p < 1 \times 10^{-18}$) (C) and normal levels of glycogen ($p = 0.44$) (D). $n > 20$ independently collected samples per value, with 25 animals per sample. Error bars represent \pm SEM. See also Figure S1.

data not shown). Notably, these include almost the entire glycolytic pathway, with genes that encode enzymes at every step in glycolysis being significantly downregulated in *dERR* mutants (Figure 2B). Northern blot hybridization was used to validate these changes in gene expression, demonstrating highly reduced transcript levels (Figure S2A). Interestingly, the *Drosophila* ortholog of *Phosphofructokinase* (*Pfk*), which




Figure 2. Genes Involved in Carbohydrate Metabolism Are Downregulated in *dERR* Mutants

(A) Gene ontology (GO) analysis of the 572 upregulated and 334 downregulated genes in *dERR*¹/*dERR*² mutant animals relative to *w*¹¹¹⁸ controls. The top GO categories for each gene set are listed in order of significance along with the number of genes affected in that category, the total number of genes in that category (in parentheses), and the statistical significance of the match.

(B) A diagram of glycolysis is depicted that displays the glycolytic genes that are downregulated in *dERR* mutants followed by their fold change in expression from the microarray.

(C) Trehalose was measured in extracts from *Pfk*^{06339/+} (Control) and *Pfk*^{06339/Df(2R)BSC303} (*Pfk*⁻) mid-second-instar larvae, revealing elevated levels in *Pfk* mutants.

(D) Trehalose concentrations were determined for *dERR*¹/*dERR*²; *da-GAL4* controls (gray box), *dERR*¹/*dERR*²; *da-GAL4* mutants (black box), and *UAS-Pgi; dERR*¹/*dERR*², *UAS-Pfk, da-GAL4* animals (white box). Trehalose levels are rescued when both transgenes are expressed with the ubiquitous *da-GAL4* driver.

Error bars represent ± SEM. See also Figure S2 and Table S1.

that these phenotypes arise from defects in other pathways (Figure S2E and data not shown).

The higher levels of trehalose seen in *dERR* mutants relative to *Pfk* mutants suggest that *dERR* plays important roles in other aspects of carbohydrate metabolism. At least one of these functions appears to be the pentose phosphate pathway (PPP), where seven of the nine genes that encode PPP enzymes are downregulated in the *dERR* mutant (Fig-

ure 2A and Tables S1 and S2). The PPP is essential for growth and energy storage, using glucose-6-phosphate to synthesize ribose-5-phosphate for nucleotide production, and generating NADPH for fatty acid synthesis and other biosynthetic reactions. *ImpL3*, which encodes lactate dehydrogenase (Ldh), is also significantly affected in *dERR* mutants, representing the seventh most highly downregulated gene (Tables S1 and S2). This enzyme converts pyruvate to lactate, regenerating NAD⁺ for use by the glycolytic pathway.

encodes the rate-limiting enzyme in glycolysis, is the fifth most downregulated gene in *dERR* mutants and the most highly affected gene in the glycolytic pathway (Table S1). These observations suggest that the elevated levels of trehalose in *dERR* mutants are due, at least in part, to decreased glycolytic flux. Consistent with this hypothesis, a mutation that eliminates *Pfk* expression results in a 48% increase in trehalose levels, demonstrating that a block in glycolysis can partially phenocopy the elevated trehalose levels observed in *dERR* mutants (Figure 2C and Figure S2). In addition, although expression of *Pfk* alone does not rescue the high trehalose levels in *dERR* mutants, expression of both *Pgi* and *Pfk* under the control of a ubiquitous GAL4 driver is sufficient to fully rescue this phenotype (Figure 2D). Moreover, ubiquitous expression of *Pgi* and *Pfk* in a *dERR* mutant background allows 8% of the animals to complete larval development and pupariate (compared with 1% of *dERR* mutant controls; *p* < 0.01), suggesting that the lethality of *dERR* mutants is due, at least in part, to the reduced expression of these two enzymes. The reduced levels of TAG and ATP, however, are not rescued in these animals, suggesting

that these phenotypes arise from defects in other pathways (Figure S2E and data not shown).

The higher levels of trehalose seen in *dERR* mutants relative to *Pfk* mutants suggest that *dERR* plays important roles in other aspects of carbohydrate metabolism. At least one of these functions appears to be the pentose phosphate pathway (PPP), where seven of the nine genes that encode PPP enzymes are downregulated in the *dERR* mutant (Fig-

Metabolomic Analysis of *dERR* Mutants

In order to characterize how these changes in gene expression might impact global metabolism, we used small-molecule gas chromatography/mass spectrometry (GC/MS) analysis to compare the metabolic profiles of *w*¹¹¹⁸ and *CantonS* controls with *dERR*¹/*dERR*² and *dERR*¹/*Df(3L)Exel6112* mutant second-instar larvae. This metabolomic approach revealed that *dERR* mutants possess increased concentrations of glucose-6-phosphate, sorbitol, mannose-6-phosphate, and three unidentified carbohydrates, consistent with their inability to metabolize




Figure 3. *dERR* Mutants Display Changes in the Levels of Specific Metabolites

(A and B) GC/MS was used to compare the relative levels of small metabolites in *CantonS* (blue) and *w*¹¹¹⁸ (green) controls with *dERR*¹/*dERR*² (orange) and *dERR*¹/*Df*(3L)*Exel6112* (red) mutant second-instar larvae. *dERR* mutants exhibit elevated levels of glucose-6-phosphate, mannose-6-phosphate, and sorbitol (A), along with diminished concentrations of lactate (B).

(C) The relative amounts of citrate, isocitrate, and succinate are similar among the four strains, whereas α -ketoglutarate, fumarate, and malate levels are decreased in mutant larvae.

(D) Methionine levels are normal in mutant animals, whereas proline concentrations are significantly lower. Glutamine and alanine levels appear to be slightly decreased in mutant strains, and aspartate is the only amino acid that is elevated in *dERR* mutants.

All data are graphically represented as a box plot, with the box representing the lower and upper quartiles, the horizontal line representing the median, and the bars representing the minimum and maximum data points. *n* = 6 samples collected from independent populations with 25 larvae per sample (see Table S3 for *p* values). Similar results were observed in two additional independent experiments (Table S3). Error bars represent \pm SEM. See also Table S3.

sugars (Figure 3A and Table S3). In contrast, oleic acid, stearic acid, and palmitic acid levels remained unchanged in mutant larvae, suggesting that fat metabolism is normal in *dERR* mutants and that the observed decrease in TAG levels are secondary to the carbohydrate defects (Table S3).

A number of metabolites downstream from glycolysis are also significantly reduced in mutant larvae. *dERR* mutants exhibit a 80%–95% decrease in lactate, which confirms that there is decreased flux through the glycolytic pathway and suggests that wild-type larvae produce a relatively large amount of lactate (Figure 3B and Table S3). Some TCA intermediates are also

reduced in *dERR* mutants, with no reproducible changes in citrate, isocitrate, or succinate levels, but a more than 90% reduction in α -ketoglutarate levels and a more than 60% reduction in fumarate and malate levels (Figure 3C and Table S3). These effects suggest that the TCA cycle is cataplerotically depleted in *dERR* mutants, with a major effect on α -ketoglutarate, which contributes to amino acid and purine biosynthesis.

Most amino acids, such as methionine, are relatively unaffected in *dERR* mutants (Figure 3D and Table S3). Interestingly, proline is the only amino acid that is reproducibly and significantly depleted in these animals. Glutamine and alanine levels are also somewhat depleted, but not consistently or to the same extent as the effects on proline (Figure 3D and Table S3). Aspartate is the only amino acid that is consistently elevated in our analysis (Figure 3D). Taken together with the depleted malate levels and reduced glycolytic flux in the *dERR* mutants, this increase in aspartate concentration may be indicative of a decrease in the activity of the malate-aspartate shuttle due to low substrate availability.

***dERR* Temporally Coordinates the Expression of Glycolytic Enzymes**

Previous studies have shown that genes involved in glycolysis are upregulated at the end of embryogenesis and that the encoded enzymes become active just prior to the onset of larval development (Wright and Shaw, 1970; Madhavan et al., 1972; Sun et al., 1988; Shaw-Lee et al., 1991, 1992; Roselli-Rehffuss et al., 1992; Currie and Sullivan, 1994a, 1994b; Abu-Shumays and Fristrom, 1997). The observation that these genes are dependent on *dERR* function raises the possibility that this receptor coordinates their expression at the end of embryogenesis, directing a metabolic switch that promotes carbohydrate metabolism. To test this hypothesis, we examined the temporal expression patterns of six *dERR* target genes during embryogenesis and early larval development in control and *dERR* mutant animals (Figures 4A and 4B). Five of these genes encode glycolytic enzymes, *Pgi* (phosphoglucose isomerase), *Pfk*, *Tpi* (triose phosphate isomerase), *Gapdh2* (glyceraldehyde 3-phosphate dehydrogenase), *Pglym78* (phosphoglycerate mutase), and *ImpL3*. Interestingly, these genes are coordinately induced at 10–14 hr after egg laying and reach maximal levels of expression just prior to larval hatching (Figure 4A). In addition, this metabolic switch is severely disrupted in *dERR* mutants (Figure 4B).

Nearly every gene that encodes a glycolytic enzyme lies near a predicted *dERR* binding site, suggesting that they are directly regulated by the receptor (Table S4). This includes *Pfk*, which has a canonical *dERR* binding site in the fourth intron that is conserved across *Drosophila* species (Figures S2B and S3A and Table S4). This sequence is bound by *dERR* in an electrophoretic mobility shift assay, and the interaction can be efficiently competed with an excess of unlabeled wild-type binding site, but not an oligonucleotide that carries a 2 bp mutation in the core binding sequence (Figure S3B). *dERR* is also bound to the *Pfk* site *in vivo*, as demonstrated by chromatin immunoprecipitation, suggesting that it represents a functional regulatory interaction (Figure S3C). To test this possibility, we examined the activity of a multimerized version of the *Pfk* *dERR* binding site placed upstream from a *lacZ* reporter gene.




Figure 4. *dERR* Is Required for the Coordinate Induction of Glycolytic Gene Expression during Mid-Embryonic Development

(A and B) Total RNA from staged *w*¹¹¹⁸ control (A) and *dERR*¹/*dERR*² mutant (B) embryos, first-instar larvae (L1), and second-instar larvae (L2) was analyzed by northern blot hybridization to detect the expression of transcripts encoding glycolytic enzymes. Glycolytic genes are coordinately induced during mid-embryogenesis in control animals (A) but not in *dERR* mutants (B). A sample of RNA from 24–28 hr *w*¹¹¹⁸ first-instar larvae (Ctrl) was included on the blot of *dERR* mutant RNA to facilitate comparisons between control and *dERR* mutant animals.

(C) The temporal expression pattern of a *lacZ* reporter construct that carries a multimerized *dERR* binding site from the *Pfk* locus was analyzed by northern blot hybridization in staged *w*¹¹¹⁸ control or *dERR*¹ mutant larvae to detect the expression of *lacZ* or *Pfk* mRNA. The reporter is expressed in synchrony with zygotic *Pfk* expression in control animals and is dependent on *dERR* function. Hybridization to detect *rp49* mRNA is included as a control for loading and transfer.

The temporal pattern of *lacZ* messenger RNA (mRNA) accumulation in transgenic embryos parallels that of the endogenous zygotic *Pfk* transcript, and this expression is almost completely abolished in a *dERR* mutant background (Figure 4C). Taken together, these results demonstrate that *dERR* can directly regulate glycolytic gene expression and that *dERR* binding is necessary and sufficient to coordinately induce these genes during mid-embryogenesis.

***dERR* Expression and Activation Are Dynamically Regulated during Mid-Embryogenesis**

The coordinate induction of *dERR* target genes in 10–142 hr embryos raises the question of how this timing is achieved. *dERR* transcripts are present throughout embryonic

development, indicating that either *dERR* protein levels and/or *dERR* activity is temporally regulated (Sullivan and Thummel, 2003). To detect *dERR* protein, we established a transformant line that carries a 3.3 kb genomic fragment spanning the *dERR* locus, with *GFP* inserted after the translation start site (*dERR-GFP*). This *dERR-GFP* reporter construct accurately reflects the activity of the native locus, as it is expressed throughout embryogenesis in the same manner as endogenous *dERR*, rescues the lethality of *dERR*¹ mutants, and restores the normal temporal expression pattern of *Pgi*, *Pfk*, and *Pglym78* in mutant embryos (Figure S4A). In spite of constant *dERR-GFP* mRNA expression, however, no GFP fluorescence is evident during the first 12 hr after egg laying (AEL) in *dERR-GFP*, *dERR*¹ embryos (Figure 5A), and *dERR-GFP* protein is not detectable on a western blot during this time (Figure S4B). Rather, GFP fluorescence is detectable and *dERR-GFP* protein begins to accumulate at 12–16 hr AEL (Figure 5 and Figure S4B), in synchrony with the coordinate induction of *dERR* target genes (Figure 4A). Expression appears to be most abundant in the muscle and epidermis (Figure 5B). This timing is consistent with the appearance of activated *dERR* LBD, as determined with the *hs-GAL4-dERR*; *UAS-GFP* ligand sensor (Palanker et al., 2006). We confirmed this result with a second *hs-GAL4-dERR*; *UAS-lacZ* reporter strain, which displayed minimal GAL4-*dERR* LBD activity during the first 12 hr of embryogenesis and became highly active in the muscle and epidermis by 12 hr AEL (Figure S5). We conclude that *dERR* is posttranscriptionally regulated and that the accumulation of activated *dERR* protein is the primary factor that drives the timing of *dERR* regulatory functions during embryogenesis.

***dERR* Regulates Carbohydrate Metabolism in Peripheral Tissues**

The *dERR* expression and activation patterns suggest that it regulates metabolism in peripheral tissues that use glucose for growth, ATP generation, and energy storage. In order to test this possibility, we used a series of tissue-specific GAL4 drivers to express a wild-type *UAS-dERR* rescue construct in a *dERR* mutant background. As described above, ubiquitous expression of *dERR* using the *da-GAL4* driver restores normal trehalose levels in mutant animals (Figure 6). Similarly, specific *UAS-dERR* expression in the fat body (*CG-GAL4* or *r4-GAL4*), muscle (*dmef2-GAL4*), or epidermis (*A58-GAL4*) significantly rescues trehalose levels in the mutant (Figure 6 and data not shown). In contrast, expression of *dERR* in the midgut (*mex-GAL4*), insulin-producing cells (IPCs) (*dilp2-GAL4*), prothoracic gland (*phm-GAL4*), corpora cardiaca (*akh-GAL4*), or Malpighian tubules (*C42-GAL4*) has no effect on the high trehalose levels (Figure 6 and data not shown).

Consistent with *dERR* regulating carbohydrate metabolism by promoting transcription of glycolytic genes, expression of the *UAS-dERR* rescue construct by the *da-GAL4* driver partially restores expression of *Pgi*, *Pfk*, and *Impl3* in a *dERR* mutant background (Figure 6). Similarly, *Pgi*, *Pfk*, and *Impl3* mRNA levels are restored in mutant animals when wild-type *dERR* is specifically expressed in the muscle or epidermis but not in the midgut or IPCs (Figure 6). *dERR*, however, appears to promote a distinct metabolic program in the fat body, with expression of wild-type *dERR* in this tissue having only minor effects on *Pfk* expression. Instead, fat body-specific expression of wild-type




Figure 5. dERR Protein Accumulation Is Temporally Regulated during Embryonic Development

Staged *dERR*¹ mutant embryos that carry a *dERR*-GFP rescue transgene were visualized by confocal microscopy to detect GFP fluorescence. No GFP was detected between 6 and 12 hr after egg laying (AEL) (A), whereas dERR-GFP accumulates in the nuclei of several cell types, including the muscle and epidermis, between 12 and 18 hr AEL (B).

dERR leads to abnormally high levels of *Pgd* mRNA, suggesting that a role for *dERR* in the fat body is to shuttle carbohydrates through the PPP (Figure 6). Finally, *dERR* may also exert a unique function in the midgut in regulating lactate metabolism. Although midgut-specific expression of wild-type *dERR* in mutant animals did not rescue the high-trehalose phenotype or restore expression of *Pgi* or *Pfk*, it did restore partial expression of *ImpL3*. Interestingly, however, the ability of tissue-specific *UAS-dERR* expression to rescue the high trehalose levels or metabolic transcriptional defects in *dERR* mutants does not correlate with its effects on viability. Expression of *UAS-dERR* in the fat body of *dERR* mutants does not restore larval viability, and only 20% of mutant animals that express *dERR* in muscle are able to complete larval development (Figure S6). In contrast, 57% of mutants survive to form pupae when *UAS-dERR* is expressed with the ubiquitous *da*-GAL4 driver (Figure S6). These results suggest that the lethality of *dERR* mutants arises from a loss of gene function in multiple tissues.

DISCUSSION

Drosophila larval development is characterized by an ~200-fold increase in body mass. This period of dramatic growth requires the efficient conversion of dietary nutrients into cellular building blocks, such as amino acids, fatty acids, and nucleotides. The metabolic basis that supports juvenile growth, however, has not been defined. The studies described here support the model that *Drosophila* larval growth depends on a form of aerobic glycolysis that is similar to the Warburg effect, and that this




Figure 6. dERR Functions in Peripheral Tissues to Control Carbohydrate Metabolism

Mid-second-instar larvae of five genotypes were tested for trehalose levels or the expression of specific *dERR* target genes: *dERR*^{1/+}; *UAS-dERR* controls (white boxes), *dERR*^{1/dERR}²; *UAS-dERR* mutants (black boxes), *dERR*^{2/+}; *GAL4* controls (blue boxes), *dERR*^{1/dERR}²; *GAL4* mutants (red boxes), or rescued *dERR*^{1/dERR}²; *UAS-dERR*, *GAL4* animals (gray boxes). Six *GAL4* transgenes were used to drive *UAS-dERR* expression: *da*-*GAL4* (Ubq, ubiquitous expression), *r4*-*GAL4* (fat body), *dmeF2*-*GAL4* (muscle), *A58*-*GAL4* (Epi, epidermis), *mex*-*GAL4* (midgut), and *dilp2*-*GAL4* (insulin-producing cells, IPCs). Total RNA was analyzed by northern blot hybridization to detect expression of *UAS-dERR*, three genes involved in glycolysis (*Pgi*, *Pfk*, and *ImpL3*), and a gene in the pentose phosphate pathway (*Pgd*). Hybridization to detect *rp49* mRNA was used as a control for loading and transfer. The apparent reduced level of *mex* > *dERR* relative to *A58* > *dERR* is an artifact of the blot hybridization. These levels of expression are comparable. In contrast, the low level of *dilp2* > *dERR* expression likely reflects the small number of cells that express the *dilp2* driver. Error bars represent ± SEM.

metabolic program is established as a mid-embryonic transcriptional switch in response to activated dERR protein.

dERR Regulates a Metabolic Program Associated with Cell Proliferation

Our microarray study of *dERR* mutant larvae revealed a profound and widespread effect on genes that control carbohydrate metabolism, including highly reduced expression of almost all genes in glycolysis and the PPP (Figures 2A and 2B and Table S1). This is consistent with the ~2-fold increase in trehalose seen in *dERR* mutants, along with highly reduced levels of ATP (Figures 1B and 1C). Mutation of the rate-limiting step in glycolysis, *Pfk*, results in elevated trehalose levels, similar to the *dERR* mutant, and ectopic expression of *Pgi* and *Pfk* in *dERR* mutants is sufficient to rescue this phenotype (Figures 2C and 2D). These studies demonstrate an essential role for *dERR* in carbohydrate catabolism during larval stages.

Metabolic profiling confirms these observations, showing significant accumulation of a number of sugars, including glucose-6-phosphate, the first intermediate in the glycolytic pathway (Figure 3 and Table S3). This study, however, also

reveals changes in the levels of a number of other key metabolites, providing a broader understanding of *dERR* function. Lactate is almost completely absent in *dERR* mutants, consistent with lactate dehydrogenase being one of the most highly downregulated genes in these animals. Levels of α -ketoglutarate, malate, and other late-stage TCA cycle intermediates are also significantly reduced in *dERR* mutant larvae, along with depletion of several amino acids, of which the most significant and reproducible is proline.

Interestingly, when taken together with the widespread effects of *dERR* on glycolysis and the PPP, these changes in metabolite levels are consistent with a form of aerobic glycolysis that is normally associated with cell proliferation (Wang et al., 1976; Vander Heiden et al., 2009). In the context of cancer, this metabolic signature is referred to as the Warburg effect (Warburg et al., 1928). The increase in carbohydrate metabolism is not designed to produce ATP, but rather promotes the synthesis of amino acids, lipids, and nucleotides, thereby supporting cellular proliferation. Both highly proliferative cells and *Drosophila* larvae shunt large quantities of glucose through the PPP, allowing them to generate ribose-5-phosphate for nucleotide synthesis and NADPH for fatty acid synthesis and other biosynthetic reactions (Geer et al., 1979; Eisenreich et al., 2004; Vander Heiden et al., 2009). Inadequate acetyl-CoA production from glycolysis in *dERR* mutants and reduced NADPH generation via the PPP likely results in decreased fatty acid synthesis, accounting for the reduced TAG levels in these animals (Figure 1E). In support of this hypothesis, TAG levels are reduced by 32% in animals that carry a loss-of-function mutation in *glucose-6-phosphate dehydrogenase*, which encodes a rate-limiting step in the PPP (data not shown, $p < 0.05$). Moreover, mutations in PPP enzymes significantly decrease the rate of carbohydrate-dependent fatty acid synthesis in larvae (Geer et al., 1979). Similarly, elevated Ldh activity, which is a hallmark of cancer metabolism, is present in normal *Drosophila* larvae, which display elevated expression of the *Drosophila* ortholog of Ldh and high levels of Ldh enzyme activity (Warburg, 1956; Rechsteiner, 1970; Abu-Shumays and Fristrom, 1997). This enzyme converts pyruvate into lactate, preventing pyruvate from entering the mitochondria and generating NAD^+ to promote maximal glycolytic flux. As a result of this diversion of pyruvate away from energy production, the TCA cycle in proliferating cells becomes dependent on amino acids derived from glutamic acid. Large amounts of glutamine are consumed by these cells to anaplerotically maintain the concentration of TCA intermediates. In an analogous manner, *Drosophila* larval metabolism appears to rely heavily on proline, which is significantly reduced in *dERR* mutants. Many insects use proline to generate energy and anaplerotically replenish the TCA cycle in the same way that proliferating yeast and cancer cells rely on glutamine (Sacktor and Childress, 1967; Arrese and Soulages, 2010). Taken together, these observations support the model that *dERR* establishes a metabolic state that is related to cellular proliferation, and that this function is essential for larval viability and growth.

Our studies also provide initial insights into the tissue-specific metabolic programs regulated by *dERR* during larval stages. When *dERR* is expressed in only the muscle or epidermis, it promotes transcription of the core glycolytic pathway (Figure 6). In contrast, specific expression of *dERR* in the fat body

upregulates *Pgd*, which encodes an essential enzyme in the PPP that is induced by sugar consumption (Cochrane et al., 1983). These results are consistent with the metabolic requirements of these different tissues. The dramatic expansion of the epidermis during larval growth, along with its production of cuticle, requires efficient glucose catabolism, as does the muscle to provide larval movement. In contrast, the fat body is one of the principle sites where sugar is processed by the PPP (Cochrane et al., 1983), and efficient lipid storage requires PPP activity (Geer et al., 1979). These observations indicate that *dERR* promotes appropriate tissue-specific metabolic programs during larval development.

dERR Triggers a Mid-Embryonic Metabolic Switch that Supports Larval Growth

A role for *dERR* in directing a metabolic state normally associated with cell proliferation provides a new context to understand how *Drosophila* larvae undergo their remarkable 200-fold increase in mass during the 4 days of larval development. It has long been known that many glycolytic enzymes are induced at the onset of larval development and that the PPP and Ldh are highly active at this stage (Wright and Shaw, 1970; Madhavan et al., 1972; Cochrane et al., 1983; Sun et al., 1988; Shaw-Lee et al., 1991, 1992; Roselli-Rehfuß et al., 1992; Currie and Sullivan, 1994a, 1994b; Abu-Shumays and Fristrom, 1997). No evidence, however, has tied these pathways together. In addition, although yeast use a similar mechanism to support their proliferation, no studies have addressed the metabolic state that accompanies juvenile development in higher organisms (Diaz-Ruiz et al., 2009). Our results suggest that *dERR* directs a coordinated metabolic program that supports the unusual growth that occurs during this stage. Our studies of the *dERR*-regulated transcriptional program also reveal that this metabolic state is established by a coordinate switch in gene expression during mid-embryogenesis (Figures 4A and 4B). Accumulation of active *dERR* protein in 12–18 hr embryos directly induces *Pfk* transcription and, probably, other key *dERR* target genes (Figure 5 and Figures S3 and S4). This transcriptional switch in turn establishes the metabolic requirements for the next stage in development. The timing of the accumulation of active *dERR* protein could be regulated at a number of levels including post-translational modifications, ligand binding, and/or cofactor recruitment. Further studies are needed to define the mechanisms that regulate this response. It is interesting to note that the use of aerobic glycolysis to support developmental growth may not be restricted to *Drosophila* larval stages. Many animals undergo exponential growth during embryonic and fetal stages of development. It will be interesting to determine whether similar metabolic states are associated with early growth in other organisms.

Implications for Mammalian ERR Function

Studies of mammalian ERR family members have largely focused on their roles in mitochondrial biogenesis and oxidative phosphorylation (for a review, see Tremblay and Giguère, 2007). Our microarray study of RNA isolated from *dERR* mutant larvae, however, identified very few genes associated with β oxidation, the TCA cycle, or the electron transport chain (Tables S1 and S2). Although genes involved in oxidoreductase activity are

upregulated in *dERR* mutants (Figure 2A), most encode cytochrome P450s with unknown functions. Similarly, a number of genes encoding mitochondrial ribosomal proteins are expressed at reduced levels in *dERR* mutants, although these effects are relatively minor (≤ 1.5 -fold). *dERR* mutant larvae have normal mitochondrial genome number and mitochondrial morphology, demonstrating no detectable effect on mitochondrial biogenesis (data not shown). These observations are consistent with a primary function for *dERR* in biomass production during larval stages, rather than energy generation and oxidative phosphorylation. Rather, we speculate that *dERR* may play a more central role in mitochondrial function during adult stages, when the fly is highly dependent on oxidative metabolism to support its increased mobility.

Conversely, several recent studies have expanded our understanding of the metabolic functions of mammalian ERR family members to include those controlled by *dERR*. *ERR γ* regulates glycolytic gene expression in the heart, *ERR α* is bound to the extended promoters of genes involved in glycolysis and the TCA cycle, and *ERR γ* regulates several genes in these pathways (Alaynick et al., 2007; Charest-Marcotte et al., 2010; Eichner et al., 2010). Similarly, *ERR α* is required for the upregulation of genes encoding glycolytic enzymes when cells are raised under hypoxic conditions (Ao et al., 2008). Our studies of *dERR* define carbohydrate metabolism as a key ancestral function for this nuclear receptor subclass, aspects of which have been conserved through evolution to mammals.

Our work also provides a new context to understand roles for mammalian ERR family members in cancer progression. Many studies have demonstrated a close association between ERR receptors and cancer (Ariazi et al., 2002; Stein et al., 2008; Yu et al., 2008). The molecular basis for this association, however, remains unclear, likely because of the functional redundancy and crosstalk between mammalian ERR paralogs. A recent paper has provided an initial step in this direction, showing that the miR-378* microRNA, which promotes the Warburg effect in BT-474 cancer cells, downregulates *ERR γ* expression in this context (Eichner et al., 2010). Our studies of the single *Drosophila* ERR family member raise the possibility that mammalian ERRs control the dramatic cellular proliferation associated with cancer through their ability to promote the Warburg effect.

EXPERIMENTAL PROCEDURES

Fly Stocks

Flies were maintained on standard Bloomington Stock Center medium with malt at 25°C. All larvae were raised on yeast paste at 25°C. The *dERR* mutant alleles were generated by ends-in homologous recombination via standard methods (Maggert et al., 2008). See the Supplemental Experimental Procedures for a detailed description. For generation of the *UAS-dERR* transgenic line, the *dERR-A* isoform was reverse transcribed and amplified from total RNA with the oligonucleotides 5'-gcaactgaataaccgatggtcg-3' and 5'-cctaactgactatattgcacctttgc-3'. The resulting complementary DNA (cDNA) was inserted into the pUAST vector and confirmed by DNA sequencing. The *Pfk-lacZ* reporter construct was generated by insertion of oligonucleotides containing tandem repeats of the *Pfk* *dERR*-binding site (5'-cctgaaggtcaccttg-3') between the EcoRI/KpnI sites and SacII/BamHI sites in the pH-Pelican vector. The *dERR-GFP* genomic rescue construct was generated by inserting the GFP coding region within the second exon of *dERR*, immediately after the start codon in *dERR-A* (see the Supplemental Experimental Procedures for

a detailed description). The *UAS-Pgi* and *UAS-Pfk* constructs were generated by amplification of the corresponding cDNAs (DGRC) with the oligonucleotide pairs (*Pgi*) 5'-gcgccgcatgcccggccactctctcc-3' and 5'-gcgccgctattactccaattggcttgatg-3' or (*Pfk*) 5'-gcgccgcatgcatcaataaaattcgagattacc-3' and 5'-gcgccgctattaggcgacggcgctcagtgctac-3' and insertion of the resulting PCR products into pUAST. All constructs were used for standard *P* element-mediated germline transformation and maintained in a *w¹¹¹⁸* background. See the Supplemental Experimental Procedures for a description of the GAL4 drivers used for tissue-specific rescue. *Df(3L)Exel6112*, which is a molecularly-defined deficiency that removes the entire *dERR* locus, was obtained from the Bloomington Stock Center.

Metabolic Analysis

Samples used for metabolic analyses were collected from independent matings on at least three different days and consisted of extracts prepared from 25 staged mid-second-instar larvae. Metabolite levels from each sample were combined to determine the average concentration and standard error, except for the GC/MS analysis, in which the three experiments were analyzed separately. Measurements of ATP, glycogen, and TAG were performed as described and normalized to total soluble protein (Palanker et al., 2009; Wang et al., 2010). For trehalose assays, 25 larvae were homogenized in trehalase buffer (5 mM Tris [pH 6.6], 137 mM NaCl, 2.7 mM KCl) and immediately incubated at 70°C for 5 min. Twenty microliters of homogenate was incubated with either 20 μ l trehalase buffer or 20 μ l trehalase buffer + 0.06 μ l porcine trehalase (Sigma, T8778-1UN) for ~12 hr at 37°C, after which the samples were centrifuged at maximum speed for 3 min. Thirty microliters of sample was transferred to a 96-well plate and incubated with 100 μ l glucose reagent (Sigma). The plate was incubated at 37°C for 30 min, after which the reaction was stopped by addition of 100 μ l 12 N H₂SO₄, and the color intensity was measured with a BioTek Synergy HT microplate reader at 540 nm. Glucose and glucose plus trehalose amounts were determined with a standard curve, and the amount of trehalose alone was determined by subtraction of the amount of glucose from the amount of glucose plus trehalose. GC/MS analysis was performed at the University of Utah Metabolomics Core Facility, as described (Wang et al., 2010).

DNA Binding

EMSA were performed essentially as described (Horner et al., 1995). The oligonucleotide containing the wild-type *dERR* binding site in *Pfk* consisted of the sequence 5'-GGCGTCTCTGGCCTGAAGTACACCTTGAAA-3' and the oligonucleotide in which the *dERR* binding site was mutated contained the sequence 5'-GGCGTCTCTGGCCTGAAGTACACCTTGAAA-3', which carries a 2 bp change in the AGGTCA core binding site. Chromatin immunoprecipitation of *dERR-GFP* was performed as described (Sieber and Thummel, 2009), with a mouse anti-GFP antibody (MBL) used to immunoprecipitate *dERR-GFP*.

Developmental Northern Blots

Staged embryos and larvae were collected as described (Sullivan and Thummel, 2003). Approximately 250 *dERR¹/dERR²* embryos were hand sorted for each time point. RNA was extracted with TriPure isolation reagent (Sigma), and northern blot analysis was conducted essentially as described (Karim and Thummel, 1991).

Microarray Analysis

Microarray analysis was performed in triplicate on independent RNA samples isolated from *w¹¹¹⁸* or *dERR^{1/2}* mid-second-instar larvae. Total RNA was extracted with TriPure (Roche) followed by purification with RNeasy columns (QIAGEN). Probe labeling, hybridization to Affymetrix GeneChip *Drosophila* Genome 2.0 Arrays, and scanning were performed by the University of Maryland Biotechnology Institute Microarray Core facility. Raw data were normalized with RMA (Bolstad et al., 2003; Irizarry et al., 2003) and analyzed with SAM, imposing a 1.5-fold cutoff and false discovery rate of $\leq 1.643\%$ (Tusher et al., 2001).

Statistical Analyses

Statistical significance was calculated with an unpaired two-tailed Student's *t* test with unequal variance. All quantitative data are reported as the mean \pm SEM.

ACCESSION NUMBERS

The NCBI GEO accession number for the microarray data reported in this paper is GSE23336.

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures, six figures, and four tables and can be found with this article online at doi:10.1016/j.cmet.2011.01.005.

ACKNOWLEDGMENTS

We thank J. Cox at the Health Sciences Center Metabolomics Core Facility at the University of Utah for assistance with the GC/MS metabolomic assays and the Bloomington Stock Center for providing fly stocks. We thank A. Sullivan for generating the *UAS-dERR* transgenic strain, H. Xie and K. Golic for help with the gene targeting, A.F. Ruaud, M. Sieber, and L. Palanker for helpful discussions and technical assistance, and M. Horner and D. Seay for comments on the manuscript. J.M.T. was supported by National Institutes of Health (NIH) National Research Service Award F32DK083864 from the National Institute of Diabetes and Digestive and Kidney Diseases. This research was supported by NIH grants 1R01DK075607 and 1RC1DK086426.

Received: July 28, 2010

Revised: November 10, 2010

Accepted: December 7, 2010

Published: February 1, 2011

REFERENCES

- Abu-Shumays, R.L., and Fristrom, J.W. (1997). IMP-L3, A 20-hydroxyecdysone-responsive gene encodes *Drosophila* lactate dehydrogenase: structural characterization and developmental studies. *Dev. Genet.* **20**, 11–22.
- Alaynick, W.A., Kondo, R.P., Xie, W., He, W., Dufour, C.R., Downes, M., Jonker, J.W., Giles, W., Naviaux, R.K., Giguère, V., and Evans, R.M. (2007). *ERR* γ directs and maintains the transition to oxidative metabolism in the postnatal heart. *Cell Metab.* **6**, 13–24.
- Ao, A., Wang, H., Kamarajugadda, S., and Lu, J. (2008). Involvement of estrogen-related receptors in transcriptional response to hypoxia and growth of solid tumors. *Proc. Natl. Acad. Sci. USA* **105**, 7821–7826.
- Ariazi, E.A., Clark, G.M., and Mertz, J.E. (2002). Estrogen-related receptor alpha and estrogen-related receptor gamma associate with unfavorable and favorable biomarkers, respectively, in human breast cancer. *Cancer Res.* **62**, 6510–6518.
- Arrese, E.L., and Soulages, J.L. (2010). Insect fat body: energy, metabolism, and regulation. *Annu. Rev. Entomol.* **55**, 207–225.
- Bolstad, B.M., Irizarry, R.A., Astrand, M., and Speed, T.P. (2003). A comparison of normalization methods for high density oligonucleotide array data based on variance and bias. *Bioinformatics* **19**, 185–193.
- Busch, B.B., Stevens, W.C., Jr., Martin, R., Ordentlich, P., Zhou, S., Sapp, D.W., Horlick, R.A., and Mohan, R. (2004). Identification of a selective inverse agonist for the orphan nuclear receptor estrogen-related receptor alpha. *J. Med. Chem.* **47**, 5593–5596.
- Chao, E.Y., Collins, J.L., Gaillard, S., Miller, A.B., Wang, L., Orband-Miller, L.A., Nolte, R.T., McDonnell, D.P., Willson, T.M., and Zuercher, W.J. (2006). Structure-guided synthesis of tamoxifen analogs with improved selectivity for the orphan *ERR* γ . *Bioorg. Med. Chem. Lett.* **16**, 821–824.
- Charest-Marcotte, A., Dufour, C.R., Wilson, B.J., Tremblay, A.M., Eichner, L.J., Arlow, D.H., Mootha, V.K., and Giguère, V. (2010). The homeobox protein *Prox1* is a negative modulator of *ERR* α /*PGC-1* α bioenergetic functions. *Genes Dev.* **24**, 537–542.
- Chawla, A., Repa, J.J., Evans, R.M., and Mangelsdorf, D.J. (2001). Nuclear receptors and lipid physiology: opening the X-files. *Science* **294**, 1866–1870.
- Cochrane, B.J., Lucchesi, J.C., and Laurie-Ahlberg, C.C. (1983). Regulation of enzyme activities in *Drosophila*: genetic variation affecting induction of glucose 6-phosphate and 6-phosphogluconate dehydrogenase in larvae. *Genetics* **105**, 601–613.
- Currie, P.D., and Sullivan, D.T. (1994a). Structure and expression of the gene encoding phosphofructokinase (PFK) in *Drosophila melanogaster*. *J. Biol. Chem.* **269**, 24679–24687.
- Currie, P.D., and Sullivan, D.T. (1994b). Structure, expression and duplication of genes which encode phosphoglyceromutase of *Drosophila melanogaster*. *Genetics* **138**, 352–363.
- Diaz-Ruiz, R., Uribe-Carvajal, S., Devin, A., and Rigoulet, M. (2009). Tumor cell energy metabolism and its common features with yeast metabolism. *Biochim. Biophys. Acta* **1796**, 252–265.
- Eichner, L.J., Perry, M.C., Dufour, C.R., Bertos, N., Park, M., St-Pierre, J., and Giguère, V. (2010). *miR-378*(*) mediates metabolic shift in breast cancer cells via the *PGC-1* β /*ERR* γ transcriptional pathway. *Cell Metab.* **12**, 352–361.
- Eisenreich, W., Ettenhuber, C., Laupitz, R., Theus, C., and Bacher, A. (2004). Isotopolog perturbation techniques for metabolic networks: metabolic recycling of nutritional glucose in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. USA* **101**, 6764–6769.
- Geer, B.W., Lindel, D.L., and Lindel, D.M. (1979). Relationship of the oxidative pentose shunt pathway to lipid synthesis in *Drosophila melanogaster*. *Biochem. Genet.* **17**, 881–895.
- Horner, M.A., Chen, T., and Thummel, C.S. (1995). Ecdysteroid regulation and DNA binding properties of *Drosophila* nuclear hormone receptor superfamily members. *Dev. Biol.* **168**, 490–502.
- Huss, J.M., Imahashi, K., Dufour, C.R., Weinheimer, C.J., Courtois, M., Kovacs, A., Giguère, V., Murphy, E., and Kelly, D.P. (2007). The nuclear receptor *ERR* α is required for the bioenergetic and functional adaptation to cardiac pressure overload. *Cell Metab.* **6**, 25–37.
- Irizarry, R.A., Bolstad, B.M., Collin, F., Cope, L.M., Hobbs, B., and Speed, T.P. (2003). Summaries of Affymetrix GeneChip probe level data. *Nucleic Acids Res.* **31**, e15.
- Karim, F.D., and Thummel, C.S. (1991). Ecdysone coordinates the timing and amounts of *E74A* and *E74B* transcription in *Drosophila*. *Genes Dev.* **5**, 1067–1079.
- Lehman, J.J., and Kelly, D.P. (2002). Transcriptional activation of energy metabolic switches in the developing and hypertrophied heart. *Clin. Exp. Pharmacol. Physiol.* **29**, 339–345.
- Luo, J., Sladek, R., Carrier, J., Bader, J.A., Richard, D., and Giguère, V. (2003). Reduced fat mass in mice lacking orphan nuclear receptor estrogen-related receptor alpha. *Mol. Cell. Biol.* **23**, 7947–7956.
- Madhavan, K., Fox, D.J., and Ursprung, H. (1972). Developmental genetics of hexokinase isozymes in *Drosophila melanogaster*. *J. Insect Physiol.* **18**, 1523–1530.
- Maggert, K.A., Gong, W.J., and Golic, K.G. (2008). Methods for Homologous Recombination in *Drosophila* (Totowa, N.J.: Humana Press).
- Ong, K.K. (2010). Early determinants of obesity. *Endocr. Dev.* **19**, 53–61.
- Palanker, L., Necakov, A.S., Sampson, H.M., Ni, R., Hu, C., Thummel, C.S., and Krause, H.M. (2006). Dynamic regulation of *Drosophila* nuclear receptor activity in vivo. *Development* **133**, 3549–3562.
- Palanker, L., Tennesen, J.M., Lam, G., and Thummel, C.S. (2009). *Drosophila* *HNF4* regulates lipid mobilization and beta-oxidation. *Cell Metab.* **9**, 228–239.
- Rechsteiner, M.C. (1970). *Drosophila* lactate dehydrogenase and alpha-glycerolphosphate dehydrogenase: distribution and change in activity during development. *J. Insect Physiol.* **16**, 1179–1192.
- Roselli-Rehffuss, L., Ye, F., Lissemore, J.L., and Sullivan, D.T. (1992). Structure and expression of the phosphoglycerate kinase (*Pgk*) gene of *Drosophila melanogaster*. *Mol. Gen. Genet.* **235**, 213–220.
- Sacktor, B., and Childress, C.C. (1967). Metabolism of proline in insect flight muscle and its significance in stimulating the oxidation of pyruvate. *Arch. Biochem. Biophys.* **120**, 583–588.
- Shaw-Lee, R.L., Lissemore, J.L., and Sullivan, D.T. (1991). Structure and expression of the triose phosphate isomerase (*Tpi*) gene of *Drosophila melanogaster*. *Mol. Gen. Genet.* **230**, 225–229.

- Shaw-Lee, R., Lissemore, J.L., Sullivan, D.T., and Tolan, D.R. (1992). Alternative splicing of fructose 1,6-bisphosphate aldolase transcripts in *Drosophila melanogaster* predicts three isozymes. *J. Biol. Chem.* **267**, 3959–3967.
- Sieber, M.H., and Thummel, C.S. (2009). The DHR96 nuclear receptor controls triacylglycerol homeostasis in *Drosophila*. *Cell Metab.* **10**, 481–490.
- Stein, R.A., Chang, C.Y., Kazmin, D.A., Way, J., Schroeder, T., Wergin, M., Dewhirst, M.W., and McDonnell, D.P. (2008). Estrogen-related receptor alpha is critical for the growth of estrogen receptor-negative breast cancer. *Cancer Res.* **68**, 8805–8812.
- Sullivan, A.A., and Thummel, C.S. (2003). Temporal profiles of nuclear receptor gene expression reveal coordinate transcriptional responses during *Drosophila* development. *Mol. Endocrinol.* **17**, 2125–2137.
- Sun, X.H., Tso, J.Y., Lis, J., and Wu, R. (1988). Differential regulation of the two glyceraldehyde-3-phosphate dehydrogenase genes during *Drosophila* development. *Mol. Cell. Biol.* **8**, 5200–5205.
- Symonds, M.E., Sebert, S.P., Hyatt, M.A., and Budge, H. (2009). Nutritional programming of the metabolic syndrome. *Nat. Rev. Endocrinol.* **5**, 604–610.
- Tremblay, A.M., and Giguère, V. (2007). The NR3B subgroup: an ovERRview. *Nucl. Recept. Signal.* **5**, e009.
- Tusher, V.G., Tibshirani, R., and Chu, G. (2001). Significance analysis of microarrays applied to the ionizing radiation response. *Proc. Natl. Acad. Sci. USA* **98**, 5116–5121.
- Vander Heiden, M.G., Cantley, L.C., and Thompson, C.B. (2009). Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science* **324**, 1029–1033.
- Wang, T., Marquardt, C., and Foker, J. (1976). Aerobic glycolysis during lymphocyte proliferation. *Nature* **261**, 702–705.
- Wang, L., Lam, G., and Thummel, C.S. (2010). Med24 and Mdh2 are required for *Drosophila* larval salivary gland cell death. *Dev. Dyn.* **239**, 954–964.
- Warburg, O. (1956). On the origin of cancer cells. *Science* **123**, 309–314.
- Warburg, O., Posener, K., and Negelein, E. (1928). Ueber den stoffwechsel der tumoren. *Biochem. Z.* **152**, 319–344.
- Willy, P.J., Murray, I.R., Qian, J., Busch, B.B., Stevens, W.C., Jr., Martin, R., Mohan, R., Zhou, S., Ordentlich, P., Wei, P., et al. (2004). Regulation of PPARgamma coactivator 1alpha (PGC-1alpha) signaling by an estrogen-related receptor alpha (ERRalpha) ligand. *Proc. Natl. Acad. Sci. USA* **101**, 8912–8917.
- Wright, D.A., and Shaw, C.R. (1970). Time of expression of genes controlling specific enzymes in *Drosophila* embryos. *Biochem. Genet.* **4**, 385–394.
- Yu, S., Wong, Y.C., Wang, X.H., Ling, M.T., Ng, C.F., Chen, S., and Chan, F.L. (2008). Orphan nuclear receptor estrogen-related receptor-beta suppresses in vitro and in vivo growth of prostate cancer cells via p21(WAF1/CIP1) induction and as a potential therapeutic target in prostate cancer. *Oncogene* **27**, 3313–3328.